

1. (a) $\text{Mg} + \text{H}_2\text{O} \rightarrow \text{MgO} + \text{H}_2$
ignore state symbols 1
 White solid / powder / ash / smoke
ignore precipitate
ignore fumes 1
 (Bright) white light / flame
allow glow
penalise effervescence under list principle 1
- (b) $2\text{Na} + \frac{1}{2}\text{O}_2 \rightarrow \text{Na}_2\text{O}$ / $4\text{Na} + \text{O}_2 \rightarrow 2\text{Na}_2\text{O}$
Allow multiples, ignore state symbols
Allow $2\text{Na} + \text{O}_2 \rightarrow \text{Na}_2\text{O}_2$ 1
 white / yellow solid / ash / smoke
ignore precipitate
ignore fumes 1
 orange / yellow flame 1
2. (a) Na_2O is an ionic lattice / giant ionic / ionic crystal
CE= 0 if molecules, atoms, metallic mentioned
Mention of electronegativity max 1 out of 2 1
 With strong forces of attraction between ions
Allow strong ionic bonds / lots of energy to separate ions 1
- (b) SO_3 is a larger molecule than SO_2
Allow greater M. / surface area 1
 So van der Waals' forces between molecules are stronger
Any mention of ions, CE= 0 1
- (c) Ionic
Do not allow ionic with covalent character 1
 Contains O^{2-} ions / oxide ions
Equations of the form $\text{O}^{2-} + \text{H}^+ \rightarrow \text{OH}^-$ / $\text{O}^{2-} + 2\text{H}^+ \rightarrow \text{H}_2\text{O}$ / $\text{O}^{2-} + \text{H}_2\text{O} \rightarrow 2\text{OH}^-$ score M2 and M3 1
 These / O^{2-} ions (accept protons to) form OH^- / hydroxide / water (must score M2 to gain M3) 1
- (d) (i) $\text{SO}_2 + \text{H}_2\text{O} \rightarrow \text{H}^+ + \text{HSO}_3^-$
Allow $2\text{H}^+ + \text{SO}_3^{2-}$ but no ions, no mark
Only score (d)(ii) if (d)(i) correct 1
 (ii) Reaction is an equilibrium / reversible reaction displaced mainly to the left / partially ionised / dissociated
Allow reaction does not go to completion 1
- (e) SiO_2 reacts with bases / NaOH / CaO / CaCO_3
Ignore incorrect formulae for silicate 1

[6]

[10]

- (b) Big difference in electronegativity leads to ionic bonding,
smaller covalent
Lose a mark if formula incorrect 1
- Sodium oxide ionic lattice 1
- Strong forces of attraction between ions 1
- P_4O_{10} covalent molecular
Must have covalent and molecular (or molecules) 1
- Weak (intermolecular) forces between molecules
Or weak vdW, or weak dipole–dipole between molecules 1
- melting point Na_2O greater than for P_4O_{10}
Or argument relating mpt to strength of forces 1
- (c) Moles $NaOH = 0.0212 \times 0.5 = 0.0106$
M1 moles of NaOH correct 1
- Moles of $H_3PO_4 = 1/3$ moles of $NaOH (= 0.00353)$
M2 is for 1/3 1
- Moles of P in 25000 l = $0.00353 \times 10^6 = 3.53 \times 10^3$
M3 is for factor of 1,000,000 1
- Moles of $P_4O_{10} = 3.53 \times 10^3/4$
M4 is for factor of 1/4 (or 1/2 if P_2O_5) 1
- Mass of $P_4O_{10} = 3.53 \times 10^3/4 \times 284 = 0.251 \times 10^6$ g
= 251 kg
(Or if P_2O_5 $3.53 \times 10^3/2 \times 142$)
M5 is for multiplying moles by M_r with correct units
allow conseq on incorrect M4
(allow 250-252) 1

[15]